


Transit Foliar®

FOLIAR-APPLIED LIQUID NUTRIENT



GUARANTEED ANALYSIS

Zinc (Zn)..... 0.5%
0.5% Chelated Zinc
 Derived from: zinc EDTA.

2.5 US Gal / 9.5 Liters
 8.7 lbs per Gallon @ 68° F
 1.0 kgs per Liter @ 20° C
 Net Weight 21.7 lbs / 9.8 kgs

DO NOT STORE BELOW 40°F (5°C)

FBSciences Inc.

153 N. Main St, Suite 100 | Collierville, TN 38017 USA

V122220 | F1837

3102043



COVER (FRONT OF LABEL) ↑

SPECIMEN LABEL

↓ BACK (BACK OF LABEL)

Transit Foliar®

DIRECTIONS FOR USE

DO NOT spray in the heat of the day or when the plant is under moisture stress. DO NOT spray to the point of runoff. Use as fine of a spray mist as possible. Allow 3-4 hours after application before sprinkler irrigating to avoid washing the product off. Shake well and/or agitate before use.

Transit Foliar should be used at the low rate when applying with additional FBS products and at the high rate when used alone or with non-FBS products. Transit Foliar mixes well with most other liquids, fertilizers, and wettable powders. A standard jar test is recommended before tank mixing. Only tank mix Transit Foliar with products where systemic uptake is desired. DO NOT mix with selective herbicides or crop oil concentrates without first consulting your local agronomist.

RATES & TIMING FOR ALL CROPS

Apply **6-15 ounces/acre** anytime during the growing season; Use higher rate with spray volume over 50 gallons/acre.

INSTRUCCIONES DE USO

NO rociar en el calor del día o cuando la planta esté bajo estrés hídrico. NO rociar hasta el punto de escurrimiento. Utilizar una niebla de rociado lo más fina posible. Dejar pasar de 3 a 4 horas después de la aplicación antes de regar con aspersión para evitar lavar el producto. Agitar bien y / o agitar antes de usar.

Transit Foliar debe usarse a una tasa baja cuando se aplica con productos de FBS adicionales y a una tasa alta cuando se usa solo o con productos que no son de FBS. Transit Foliar se mezcla bien con la mayoría de los demás líquidos, fertilizantes y polvos humectables. Se recomienda una prueba de jarra estándar antes de mezclar en el tanque. Solo mezclar en el tanque Transit Foliar con productos donde se desee una absorción sistémica. NO mezclar con herbicidas selectivos o concentrados de aceite de cultivo sin antes consultar a su agrónomo local.

TASAS Y MOMENTO PARA TODOS LOS CULTIVOS

Aplicar de **6 a 15 onzas/acre** en cualquier momento durante la temporada de crecimiento; Usar una dosis más alta con un volumen de rociado superior a 50 galones/acre.

WARNING

READ ENTIRE LABEL BEFORE USE | KEEP OUT OF REACH OF CHILDREN



HAZARD STATEMENTS Harmful if swallowed. May be harmful in contact with skin.

AVISO PELIGRO Nocivo si se ingiere. Puede ser nocivo en contacto con la piel.

PRECAUTIONARY STATEMENTS Wear protective gloves/protective clothing/eye protection. Wash skin thoroughly after handling. Do not eat, drink or smoke when using this product. **IF SWALLOWED:** Rinse mouth. Call a POISON CENTER or doctor/physician if you feel unwell. Dispose of contents/container in accordance with local/national regulations.

WARRANTY AND CONDITIONS OF SALE Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. BEFORE OPENING, see other conditions and limitations that apply which are available at: <http://www.FBSciences.com/ProductConditionsOfSale/>. See patent and other patents pending: <http://www.fbsciencespatents.com>. Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.aapfco.org/metals.html>